

Title	冷感剤メントールによって誘起される膜の物理化学特性変化に関する研究
Author(s)	Gusain, Pooja
Citation	
Issue Date	2017-06
Type	Thesis or Dissertation
Text version	ETD
URL	http://hdl.handle.net/10119/14751
Rights	
Description	Supervisor:高木 昌宏, マテリアルサイエンス研究科, 博士

Detailed Membrane Physiochemical Pathways Involved in Cold-sensitization Induced by l-menthol


Takagi Laboratory
s1440152
Pooja Gusain

Background: Lipid rafts are specialized membrane microdomain enriched in saturated lipid, sphingolipids, and cholesterol¹. These regions are rather tightly packed in the liquid ordered phase, unlike the disordered phase present in the bulk of the biological membrane. Lipid rafts serve as a platform for the regulation of various cellular processes involved in signal transduction and membrane trafficking². Some membrane proteins, such as ion channels, are localized on lipid rafts. The activity of a channel is controlled by specific molecules, and changes in activity resulting from molecular binding in a channel generate a sensation, such as warmth, cold, or anaesthesia. This binding molecule may not only affect the channel, but also the lipid bilayer, thus altering the physical properties of the membrane. Therefore, channel activity is also influenced through the lipid membrane, and specific changes in the physical properties of lipid membranes by several sensing molecules have been revealed³. The two important parameters: hydrophobic acyl chain ordering and hydrophilic head group interaction could represent the sensing mechanism leading to channel activation. Hence, it becomes very important to perceive these structural changes of lipid on interaction with such sensing molecules.

The complexity of living system makes understanding the importance of chirality challenging. Biological molecules are often chiral, such as DNA, proteins, carbohydrates, lipids, and steroids. Menthol is also chiral. There are several diastereomers of menthol, of which the most common are (1S, 2R, 5S)-2-isopropyl-5-methylcyclohexanol (d-menthol) and (1R, 2S, 5R)-2-isopropyl-5-methylcyclohexanol (l-menthol). l-Menthol is the main form found in nature and is widely used in products such as toothpaste, chewing gum, and cigarettes. l-Menthol is believed to bind onto TRPM8 and causes a well-known, desired, cooling sensation⁴. In contrast, d-menthol does not exist in nature and does not induce cooling sensation. The underlying significant differences in the interaction between d-/l-menthol and the lipid membrane are not well understood. The first step in addressing this is to elucidate how and where d-/l-menthol each interact with a membrane. Considering these facts, we used GUVs as model membranes (artificial cells) and live cells to examine the interaction between a lipid membrane and d-/l-menthol.

Research Objective: Each part of the human body acts for the original roles and every role for each sense is far more crucial for their proper functioning. One of the important functions of our body is the sensing various external stimuli. The different sensitization includes cold, warm, anesthesia, etc. which is provoked by activation of some sensing channels. The activity of these channels is induced by external molecules leads to opening/closing of the channels, thus changes the physiological states of the membrane. Menthol, a chiral molecule, is one of the candidates, inducing cold sensitization to the body by activating a sub family of ion channels, termed as transient receptor potential melastatin 8 (TRPM8). Although many studies have been reported about the interaction between menthol and membrane, the mechanism at the molecular level is still remains unclear. With this understanding, I aimed to scratch physiological changes induced by menthol on the model membrane and related mechanism for its cooling behaviour.

Experimental: To study the detailed mechanistic pathways at a fundamental level, I worked on both cell-sized liposomes, i.e., GUVs as a model membrane and biological membrane specifically HEK293 cell line. By using both model membrane and biological membrane, I have shown the different localization and interaction sites of d- and l-menthol. In contrast, the role of cholesterol to alter the physiochemical properties of membrane was also demonstrated. Preparation of liposomes and staining protocol for HEK293 cells are depicted in scheme 1. The detailed procedure for the sample preparations have been discussed in chapter 2 and 5.



Scheme 1. Preparation of Liposomes (a) and staining protocol for HEK293 cells (b)

Results and discussion: In chapter 1, first we studied the initial interaction of menthol on homogeneous model membrane composed of DOPC lipid, Cholesterol, and l-menthol. We have shown that l-menthol can interact directly with the model membrane and has specific interaction. Figure 1 (a, b). Shows typical phase-contrast microscopic images in both DOPC/Chol (control) system and DOPC/Chol/menthol. The vesicle consisting of DOPC/Chol retained the spherical shape, even if the temperature was increased from 21 °C to 25 °C. On the other hand, the membrane fluctuation was observed in DOPC/Chol/menthol with same temperature change. Previous studies showed that menthol has dual behaviours, at higher temperature and concentration it has a tendency to exhibit itching/burning sense while at low temperature and concentration has cool sense. From fluctuation data, we could observe the clear concentration effect of menthol on the model membrane. The effect of menthol concentration on the thermo-sensitive fluctuation of DOPC/Chol liposomes is depicted in Figure 1(c).




Fig.1 Typical microscopic images (a, b). Thermal-responsiveness response of menthol-containing lipid vesicles (c). Percentage of lipid vesicles, which started fluctuating at a given level of temperature increase. The vesicles contained DOPC/Chol (black), 2.5 % menthol (light grey dashed), 5 % menthol (blue), 10 % menthol (dark grey dashed), 20 % menthol (black dotted), and 30 % (grey solid line). (n=30).

At lower concentration of menthol, the maximum fluctuation could be observed, while at a higher concentration rate of fluctuation declined. Generally, the thermo-induced membrane fluctuation was caused by the acquisition of the excess membrane area. The absence of fluctuation in the DOPC/Chol vesicles suggested that the membrane area remained steady. Dynamic real-time observation of membrane dynamics revealed that the menthol-containing membrane was more thermo-responsive than without menthol. The increase in the temperature enhanced the number of water molecules near polar head groups, leading to an increase in steric repulsive interaction between them and resulted into the excess surface area of liposomes. As a consequence, liposomes started fluctuating depending on the hydrophilicity of the head group. Further, to elaborate more

about the specific interaction of menthol involved in cold-sensitization, we examined the chirality-dependent interaction in both homogeneous and heterogeneous systems discussed in chapter 3 and 4.

In order to understand the effects of d- or l-menthol on the phase separation more clearly, the miscibility temperature was observed over the temperature range from 18 °C to 38 °C using thermo-controller. First, we observed the phase separation of binary lipid mixtures consisting of the unsaturated lipid DOPC and the saturated lipid DPPC without Chol at temperature ranging from 18 °C to 38 °C (obtained using thermo controller). The miscibility temperature T_m is defined as the temperature at which 50 % of GUVs become heterogeneous. Figure 2 shows the fraction of liposomes forming phase-separated structures at each temperature tested. No clear difference in miscibility temperature was obtained by adding d- or l-menthol to the binary system DOPC/DPPC, as shown in Fig. 2(a) and (d). Next, we observed phase separation in the DOPC/DPPC/Chol (Chol = 20 %) ternary lipid mixture. Also, the system without menthol and the d-/l-menthol-containing systems showed no significant differences in miscibility temperature, as shown in Fig. 2(b) and (d). At Chol = 30 %, however, clear and significant phase behaviour differences induced by d- or l-menthol were apparent, as shown in Fig. 2(c) and (d). Notably, d-menthol stabilized phase separation, whereas l-menthol dramatically lowered the phase-separated fraction. These findings can be explained by considering the change in the lipid bilayer physical state near phase transition. From the results obtained we proposed a model showing possible interaction of d-/l-menthol with lipid bilayer which could affect the cold sensing properties of TRPM8. We also believe that TRP channels can sense the temperature-dependant changes in the lipid bilayer.




Fig. 2 Miscibility temperature measurement for DOPC/DPPC/Chol/menthol system at Chol = 0 % in (a), 20 % in (b), and 30 % in (c). Black lines and black squares indicate DOPC/DPPC/Chol without menthol. Red lines with red circles and blue lines with blue triangles are DOPC/DPPC/Chol with d-menthol and l-menthol, respectively. Change in the miscibility temperature as a function of Chol concentration (d).

Conclusions: In current issue, I have explored deep understanding of the membrane interaction and physiochemical changes upon external stimuli *i.e.*, menthol at the molecular level. Moreover, the mechanical pathway involved in the cold-sensitization induced by l-menthol was clarified. I successfully demonstrated first attempt to assess the interaction of signalling molecule “menthol” with model cell membrane. Different approaches had been attempted to clarify physiological changes and morphology change induced by menthol along with the importance of chirality in the biological processes. These dynamic changes in the membrane properties enable to picture the related mechanism behind sensation such as warmth, cold, and anaesthesia. Therefore, studies on lipid rafts and membrane lipids could offer more information to unravel the mechanism involved in the channel functioning.

References:

- 1) Simons, K. & Toomre, D. Lipid rafts and signal transduction. *Nat Rev Mol Cell Biol* **1**, 31–39 (2000).
- 2) van Meer, G., Voelker, D. R. & Feigenson, G. W. Membrane lipids: where they are and how they behave. *Nat Rev Mol Cell Biol* **9**, 112–124 (2008).
- 3) Kopeć, W., Telenius, J. & Khandelia, H. Molecular dynamics simulations of the interactions of medicinal plant extracts and drugs with lipid bilayer membranes. *FEBS J.* **280**, 2785–2805 (2013).
- 4) BG, G. Menthol modulates oral sensations of warmth and cold. *Physiol Behav* **35**, 427–34 (1985).

Keywords : Lipid rafts, d- and l-menthol, membrane fluctuation, phase separation, miscibility temperature